

Methyl Pothoscandensate, a New *ent*-18(4 → 3)-Abeokaurane from *Pothos scandens*

by Hong-Xin Liu^{a)}), Jun-Long Bi^{c)}, Yue-Hu Wang^{a)}, Wei Gu^{a)}), Yao Su^{d)}, Fang Liu^{d)}, Shi-Xian Yang^{d)}, Guang-Wan Hu^{a)}, Ji-Feng Luo^{a)}, Ge-Fen Yin^{*c)}, and Chun-Lin Long^{*a)})

^{a)} Key Laboratory of Economic Plants and Biotechnology, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, P. R. China (phone/fax: + 86-10-68930381; e-mail: long@mail.kib.ac.cn)

^{b)} Graduate University of Chinese Academy of Sciences, Beijing 100049, P. R. China

^{c)} College of Animal Science and Technology, Yunnan Agricultural University, Kunming 650201, P. R. China (e-mail: yingefen383@sohu.com)

^{d)} College of Horticulture and Landscape, Yunnan Agricultural University, Kunming 650201, P. R. China

^{e)} College of Life and Environmental Sciences, Minzu University of China, Beijing 100081, P. R. China

Methyl pothoscandensate (**1**), a new molecular skeleton of *ent*-18(4 → 3)-abeokaurane, along with eight known compounds was isolated from the whole plant of *Pothos scandens*. The structure of the new compound was established by spectroscopic techniques and confirmed by single-crystal X-ray diffraction. The inhibitory activity of selected compounds against porcine respiratory and reproductive syndrome virus (PRRSV) was measured by the cytopathic effect (CPE) method. Compound **1** showed weak effect on PRRSV with an IC_{50} value of $40.3 \pm 8.3 \mu\text{M}$ ($TI = 15.7$).

Introduction. – The genus *Pothos* is a medicinally important member in the plant family Araceae, consisting of *ca.* 75 species. There are five naturally occurring species found in China, mainly distributed in the south and southwest of Yunnan Province [1]. Some species of this genus have been used in treatment of traumatic injuries, fractures, and inflammation in Chinese traditional medicinal systems. Furthermore, boiled water decoctions of *P. scandens* leaves are used as tea by the Dai people [2]. Though this genus includes many individuals used as medicinal herbs, there have been very few chemical investigations on *Pothos*. As a result, we conducted the investigation of the chemical constituents of *P. scandens*. Repeated column chromatography and recrystallization of the AcOEt extract of the whole plant yielded a novel diterpenoid **1** (Fig. 1), as well as eight known compounds, *N*-*trans*-cinnamoyltyramine (= (2*E*)-*N*-[2-(4-hydroxyphenyl)ethyl]-3-phenylprop-2-enamide) [3], *N*-*trans*-feruloyltyramine (= (2*E*)-3-(4-hydroxy-3-methoxyphenyl)-*N*-[2-(4-hydroxyphenyl)ethyl]prop-2-enamide) [4], *N*-*trans*-*p*-cumaroyltyramine (= (2*E*)-3-(4-hydroxyphenyl)-*N*-[2-(4-hydroxyphenyl)ethyl]prop-2-enamide) [5], (–)-serotobenine (= *rel*-(2*R*,2*aR*)-2,2*a*,4,5,6,8-hexahydro-2-(4-hydroxy-3-methoxyphenyl)-3*H*-furo[2,3,4-*kl*]pyrrolo[4,3,2-*fg*][3]bentatocin-3-one) [6], (3β)-*ent*-kaurane-3,16,17-triol (**2**) [7], (+)-syringaresinol (= 4,4'-(tetrahydro-1*H*,3*H*-furo[3,4-*c*]furan-1,4-diyl)bis[2,6-dimethoxyphenyl]) [8], (3β)-*ent*-kaurane-3,16,17-triol 3-β-*D*-glucopyranoside) [9], and (2*R*)-2-hydroxy-2-phenylacetonitrile 2-[*O*-β-*D*-xylopyranosyl-(1 → 6)-β-*D*-glucopyranoside] [10]. Herein we report the isolation and structure elucidation of **1**, and the results of the bioassay.




Fig. 1. *Methyl pothoscandensate (1)*, isolated from *Pothos scandens*

Results and Discussion. – Compound **1** was isolated as colorless needle crystals and its molecular formula was established as $C_{21}H_{32}O_4$ based on the HR-ESI-MS (m/z 349.2381 ($[M+H]^+$, calc. 349.2378)), which requires 6 degrees of unsaturation. The ^{13}C -NMR and DEPT spectrum of **1** (Table) showed 21 C-atom signals including those of three Me, nine CH_2 , and three CH groups and six quaternary C-atoms, of which the signals of an ester C-atom ($\delta(C)$ 170.2), two olefine C-atoms ($\delta(C)$ 145.4 and 123.7), two O-bearing C-atoms ($\delta(C)$ 81.8 and 66.3), and one MeO group ($\delta(C)$ 51.2) were detected. By comparing the NMR data with those of (3 β)-*ent*-kaurane-3,16,17-triol (**2**; see Scheme below) [7], **1** was determined to be an *ent*-kaurane-type compound. Analysis of $^1H,^1H$ -COSY data of **1** (Fig. 2) established the two segments $CH_2(1)CH_2(2)$ (**a**) and $CH_2(12)CH(13)CH_2(14)$ (**b**) (Fig. 2). In the HMBC spectrum, the correlations $CH_2(14)/C(9)$ ($\delta(C)$ 53.1), $C(12)$ ($\delta(C)$ 25.9), $C(15)$ ($\delta(C)$ 53.0), and $C(16)$ ($\delta(C)$ 81.8), $CH_2(11)/C(8)$ and $C(13)$, and $H-C(13)/(11)$, revealed the presence of a bicyclo[3.2.1]octane system (rings C and D). Based on the HMBCs $CH_2(1)/C(20)$ and $C(3)$, $CH_2(2)/C(10)$, $H-C(5)/C(1)$, $CH_2(6)/C(8)$ and $C(10)$, $CH_2(7)/C(15)$, $H-C(9)$ to $C(1)$ and $C(5)$, $Me(19)/C(3)$, $C(4)$, and $C(5)$, and $Me(20)/C(1)$, $C(5)$, $C(9)$, and $C(10)$, rings A and B were assigned. The two OH groups were located at

Table. 1H - and ^{13}C -NMR Data (400 und 100 MHz; $CDCl_3$) of Compound **1**. δ in ppm, J in Hz.

Position	$\delta(H)$	$\delta(C)$	Position	$\delta(H)$	$\delta(C)$
$CH_2(1)$	1.86–1.89, 1.05–1.08 (2m)	35.6 (CH_2)	$CH_2(12)$	1.62–1.65, 1.54–1.57 (2m)	25.9 (CH_2)
$CH_2(2)$	2.32–2.36 (m)	24.2 (CH_2)	$H-C(13)$	2.07 (br. s)	45.3 (CH)
$C(3)$		123.7 (C)	$H_\alpha-C(14)$	2.01 (d, $J=11.9$)	37.6 (CH_2)
			$H_\beta-C(14)$	1.63–1.66 (m)	
$C(4)$		145.4 (C)	$CH_2(15)$	1.58–1.61, 1.48–1.51 (2m)	53.0 (CH_2)
$H-C(5)$	1.86–1.88 (m)	50.2 (CH)	$C(16)$		81.8 (C)
$CH_2(6)$	1.86–1.89, 1.39–1.42 (2m)	22.0 (CH_2)	$CH_2(17)$	3.81, 3.69 (2d, $J=10.9$)	66.3 (CH_2)
$CH_2(7)$	1.67–1.69, 1.54–1.57 (2m)	41.0 (CH_2)	$C(18)$		170.2 (C)
$C(8)$		44.1 (C)	$Me(19)$	1.92 (s)	18.1 (Me)
$H-C(9)$	1.05–1.08 (m)	53.1 (CH)	$Me(20)$	0.89 (s)	14.3 (Me)
$C(10)$		37.1 (C)	(MeO)	3.71 (s)	51.2 (Me)
$CH_2(11)$	1.66–1.69, 1.56–1.59 (2m)	18.5 (CH_2)			




Fig. 2. $^1H,^1H$ -COSY (—) and key HMBC ($H \rightarrow C$) features of **1**

C(16) and C(17), respectively, by the correlations $\text{CH}_2(14)/\text{C}(16)$, $\text{H}-\text{C}(13)/\text{C}(11)$ and $\text{CH}_2(17)/\text{C}(13)$ and C(15). Although the correlations between $\text{CH}_2(2)$ and C(18) were not observed in the HMBC spectrum, the methoxycarbonyl group should be at the quaternary C-atom (C(3)). At last, the planar structure of **1** was deduced as shown in Fig. 2. The partial relative configuration of **1** was deduced by the analysis of ROESY correlations (Fig. 3). The key ROESY correlations $\text{H}-\text{C}(5)/\text{H}-\text{C}(9)$ showed that the two H-atoms were cofacial, and were arbitrarily assigned to be β -oriented, while Me(20) was assigned the α -orientation. The correlation of $\text{CH}_2(14)/\text{Me}(20)$ and $\text{H}-\text{C}(13)/\text{CH}_2(14)$ confirmed $\text{H}-\text{C}(13)$ to be α -oriented. The entire structure of **1** was finally determined by a single-crystal X-ray diffraction analysis (Fig. 4) [11]. Thus, **1** was elucidated as 16,17-dihydroxy-18(4 \rightarrow 3)-abeo-*ent*-kaur-3-en-18-oic acid methyl ester and given the trivial name methyl pothoscandensate.




Fig. 3. Key ROESY ($\text{H} \leftrightarrow \text{H}$) correlations of **1**





Fig. 4. ORTEP Drawing of the X-ray structure of **1**

A possible biogenetic pathway of **1** is proposed as shown in the Scheme. Compound **2** would be transformed into **iii** by Wagner–Meerwein rearrangements. Firstly, the OH group would leave from **2** to generate the cation intermediate **i** under acidic conditions.

Afterwards, one of the Me groups at C(4) would migrate to C(3) forming the intermediate **ii**, followed by H-atom elimination to give the key intermediate **iii**. Then, **iii** could be finally transformed to **1** by selectively oxidizing the allylic Me group to the acid followed by esterification.

Scheme. Proposed Biogenetic Pathway of **1**



The inhibitory activity of methyl pothoscandensate (**1**), *N-trans*-feruloyltyramine, *N-trans-p*-cumaroyltyramine, (–)-serotobenine, (3 β)-*ent*-kaurane-3,16,17-triol (**2**), (+)-syringaresinol, and (2*R*)-2-hydroxy-2-phenylacetonitrile 2-[O- β -D-xylopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside] against porcine respiratory and reproductive syndrome virus (PRRSV) was measured by the cytopathic effect (CPE) method [12]. Compound **1** showed a weak inhibitory effect on PRRSV with an IC_{50} value of $40.3 \pm 8.3 \mu\text{M}$ (TI (therapeutic index) = 15.7) compared with tilmicosin phosphate ($IC_{50} = 225.1 \pm 27.4 \mu\text{M}$, $TI = 3.8$). The other tested compounds were inactive ($IC_{50} > 200 \mu\text{M}$). Furthermore, by the real-time fluorescent quantitative reverse transcription-polymerase chain reaction (FQ RT-PCR) [13–15], the relative expression ratio of PRRSV ORF 7 and NSP9 genes was tested. ORF 7 and NSP9 mRNA relative expression level was significantly reduced by compound **1** at the concentration of 100 μM or more ($P < 0.001$; Fig. 5).




Fig. 5. ORF7 (left) and NSP9 (right) mRNA relative expression level reduced by compound **1** (* $P < 0.001$)

This work was funded by the National Natural Science Foundation of China (Grant Nos. 20972166, 31070288, 30860205, and 31161140345) and the Ministry of Education (Grant Nos. B08044 and

MUC985–9). We appreciate Dr. *Xiao-Nian Li*, Kunming Institute of Botany, for the measure and elucidation of the crystal structure. Special thanks go to *Adam Negrin*, City University of New York (Lehman College), for his reading and editing of the English.

Experimental Part

General. Column chromatography (CC): *MCI* gel (70–150 μm ; *Mitsubishi Chemical Corporation*), *C₁₈* silica gel (SiO_2 , 40–75 μm ; *Fuji Silysia Chemical Ltd.*), *Sephadex LH-20* gel (*GE Healthcare Bio-Sciences AB*), and silica gel (SiO_2 , 80–100, 20–300, and 300–400 mesh; *Qingdao Meigao Chemical Co.*). TLC: pre-coated SiO_2 *F₂₅₄* plates (*Qingdao Meigao Chemical Co.*); spots were detected under UV light (254 and 365 nm), and by spraying with 5% aq. H_2SO_4 in EtOH, followed by heating. M.p.: *X-4* melting-point apparatus (*Yingyu Yuhua Apparatus Factory*, Gongyi, P. R. China). Optical rotations: *Jasco-DIP-370* automatic digital polarimeter. UV Spectra: *Shimadzu-210A* double-beam spectrometer; λ_{max} (log ϵ) in nm. IR Spectra: *Bio-Rad-FTS-135* spectrophotometer; KBr pellets; ν in cm^{-1} . NMR Spectra: *Bruker-AM-400* and *-DRX-500* spectrometers; δ in ppm rel. to Me_4Si as internal standard, *J* in Hz. ESI-MS and HR-ESI-MS: *API-Qstar-Pulsar-1* instrument; in *m/z*.

Plant Material. The whole plant of *Pothos scandens* (8.3 kg) was collected from Xishuangbanna of Yunnan Province, P. R. China, in October 2010, and identified by Dr. *Guang-Wan Hu*, Kunming Institute of Botany. A voucher specimen (No. LHX-0091) was deposited with the Key Laboratory of Economic Plants and Biotechnology, Kunming Institute of Botany.

Extraction and Isolation. The air-dried powder of the plant material (8.3 kg) was exhaustively extracted with MeOH a total of 3 times, and the extract (0.79 kg) was suspended in H_2O and partitioned into three fractions with petroleum ether (*A*, 100 g), AcOEt (*B*, 165 g), and H_2O (*C*, 200 g). Subsequently, *Fr. B* was subjected to CC (SiO_2 $\text{CHCl}_3/\text{MeOH}$ 1:0 \rightarrow 0:1): *Frs. 1–6*. *Fr. 2* ($\text{CHCl}_3/\text{MeOH}$ 15:1) was separated by CC (*RP-18*, $\text{MeOH}/\text{H}_2\text{O}$ 2:8 \rightarrow 9:1): *Frs. 2.1–2.3*. *Fr. 2.1* was fractionated by CC (*Sephadex LH-20*, MeOH; SiO_2 , $\text{CHCl}_3/\text{MeOH}$ 10:1): **2** (40.5 mg). *Fr. 2.2* was fractionated by CC (*Sephadex LH-20*, MeOH; SiO_2 , $\text{CHCl}_3/\text{MeOH}$ 15:1): (+)-syringaresinol (13.4 mg). *Fr. 3* ($\text{CHCl}_3/\text{MeOH}$ 10:1) was separated by CC (*RP-18*, $\text{MeOH}/\text{H}_2\text{O}$ 3:7 \rightarrow 9:1): *Frs. 3.1–3.6*. *Fr. 3.1* was fractionated by CC (*Sephadex LH-20*, MeOH; silica gel, $\text{CHCl}_3/\text{MeOH}$ 10:1): (–)-serotobenine (11.0 mg). *Fr. 3.2* was fractionated by CC (*Sephadex LH-20*, MeOH; SiO_2 , petroleum ether/ Me_2CO 3:1, $\text{CHCl}_3/\text{MeOH}$ 10:1): N-trans-feruloyltyramine (29.0 mg) and N-trans-p-cumaroyletyramine (113.0 mg). *Fr. 3.3* was fractionated by CC (*Sephadex LH-20*, MeOH; SiO_2 , $\text{CHCl}_3/\text{MeOH}$ 10:1): N-trans-cinnamoyltyramine (4.2 mg). *Fr. 3.4* was fractionated by CC (*Sephadex LH-20*, MeOH; SiO_2 , petroleum ether/ Me_2CO 5:1): **1** (64.5 mg). *Fr. 5* ($\text{CHCl}_3/\text{MeOH}$ 3:1) was separated by CC (*RP-18*, $\text{MeOH}/\text{H}_2\text{O}$ 6:4 \rightarrow 9:1): *Frs. 5.1–5.3*. *Fr. 5.2* was fractionated by CC (*Sephadex LH-20*, MeOH; SiO_2 , $\text{CHCl}_3/\text{MeOH}$ 5:1): (2*R*)-2-hydroxy-2-phenylacetone nitrile 2-[*O*- β -D-xylopyranosyl-(1 \rightarrow 6)- β -D-glucopyranoside] (15.0 mg). *Fr. 5.3* was fractionated by CC (*Sephadex LH-20*, MeOH; SiO_2 , $\text{CHCl}_3/\text{MeOH}$ 4:1): (3 β)-ent-kaurane-3,16,17-triol 3-(β -D-glucopyranoside) (32.0 mg).

X-Ray Crystallographic Analysis of **1.** $\text{C}_{42}\text{H}_{64}\text{O}_8$ ($2 \times \text{C}_{21}\text{H}_{32}\text{O}_4$), M_r 696.93, colorless needle crystal, size $0.03 \times 0.12 \times 0.78$ mm³, monoclinic, space group *P2₁*; $a = 9.319(7)$, $b = 7.234(6)$, $c = 27.87(2)$ Å, $\alpha = \gamma = 90.00$, $\beta = 90.180(12)^\circ$, $V = 1878(3)$ Å³, T 296(2) K, $Z = 2$, $\rho_{\text{calc.}} = 1.232$ g/cm³; $F(000) = 760$, 18137 reflections in $-12 \leq h \leq 12$, $-9 \leq k \leq 9$, $-37 \leq l \leq 7$, measured in the range $0.73^\circ \leq \theta \leq 28.97^\circ$, completeness $\theta_{\text{max}} = 94.7\%$, g.o.f. = 0.832. Final *R* indices: $R_1 = 0.1257$ and $wR_2 = 0.2601$, Flack parameter $-2(4)$, largest difference peak and hole = 0.347 and -0.297 e Å⁻³. The intensity data for **1** were collected on a *Bruker APEX DUO* diffractometer with graphite-monochromated MoK_α radiation. The structure of **1** was solved by direct methods (SHELXS97 [11]), expanded with difference Fourier techniques, and refined by the program and full-matrix least-squares calculations. The non-H-atoms were refined anisotropically, and H-atoms were fixed at calculated positions. CCDC-848942 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via http://www.ccdc.cam.ac.uk/data_request/cif.

Methyl Pothoscandensate (=16,17-Dihydroxy-18(4 \rightarrow 3)-abeo-ent-kaur-3-en-18-oic Acid Methyl Ester = Methyl ent-16,17-Dihydroxy-18-norkaur-3-ene-3-carboxylate; **1**): Colorless needles (MeOH).

M.p. 102–103°. $[\alpha]_{\text{D}}^{23.4} = -103.5$ ($c = 0.20$, CHCl_3). UV (CHCl_3): 241.4 (3.12). IR: 3424, 1712, 1620. ^1H - and ^{13}C -NMR: Table. ESI-MS (pos.): 371 ($[\text{M} + \text{Na}]^+$). HR-ESI-MS (pos.): 349.2381 ($[\text{M} + \text{H}]^+$, $\text{C}_{21}\text{H}_{33}\text{O}_4^+$; calc. 349.2378).

Cytopathic Effect Inhibition Assay. The 50% cytotoxic concentration (CC_{50}) of compounds **1**, **2** *N*-trans-feruloyltyramine, *N*-trans-*p*-cumaroyltyramine, (–)-serotobenine, (+)-syringaresinol, (2*R*)-2-hydroxy-2-phenylacetonitrile 2- $[\text{O}-\beta\text{-D-xylopyranosyl-(1} \rightarrow 6)\text{-}\beta\text{-D-glucopyranoside}]$ and tilmicotin phosphate (positive control; Hubei Hengshuo Chemical Co., Ltd., China) against Marc-145 cells (Cell Bank of the Chinese Academy of Sciences, Shanghai) was measured according to previously described protocols [16–18]. The antiviral activity of tested compounds against PRRSV was evaluated by the cytopathic effect (CPE) inhibition assay [12]. The YN-1 strain of PRRSV was isolated from local pigs in Yunnan Province, P. R. China [19]. The tissue culture medium infective dose (TCID_{50}) of 500 viral particles with twofold serial dilutions of the compounds were added to each test well, and the plates were re-incubated for 4 d to allow development of a cytopathologic effect (CPE) if any. A noninfection control was made in the absence of natural products, and tilmicotin phosphate was used for drug control. The concentration reducing CPE by 50% with respect to virus control was estimated from graphic plots and was defined as 50% inhibited concentration (IC_{50}). The therapeutic index (*TI*) was calculated from the ratio $\text{CC}_{50}/\text{IC}_{50}$.

PRRSV mRNA Expression Inhibition Assay. The mRNA expression of PRRSV ORF7 and NSP9 genes was determined by real-time RT-PCR [13–15]. Briefly, after 4 d of incubation, total virus RNA of both administration and control groups was isolated by means of *RNAiso*TM Plus (*TaKaRa* Biotechnology, Dalian, P. R. China), dissolved in 30 μl of RNase-free H_2O (*TaKaRa*), and then stored at -80° . According to the GenBank data base, accession No. PRU87392, primers were selected and designed from conserved regions based on the ORF7 and NSP9 sequences by using Primer5.0 and Oligo6.0 software. A 330 base pair fragment of the PRRSV ORF7 gene was amplified by using the following primers: forward primer was 5'-AATGGCCAGCCAGTCAATCA-3' and reverse primer was 5'-TCATGCTGAGGGTGATGCTG-3'. A 162 base pair fragment of the PRRSV NSP9 gene was amplified by using the following primers: forward primer was 5'-CACTAAAGAGGAAGTCGCACTCA-3' and reverse primer was 5'-GGTATGTCTCCAAACCTGTATTCTG-3'. A 130 base pair fragment of the beta-actin gene was amplified by using the following primers: forward primer was 5'-ATCCAGGCTGTGCTGTCC-3' and reverse primer was 5'-GAGGATCTTCATGAGGTAGTCG-3'.

cDNAs were synthesized with the *PrimeScript*[®] RT reagent kit (*TaKaRa*) with 10 μl of reaction mixtures containing 4.5 μl of RNase free dH_2O , 2 μl of $5 \times \text{PrimeScript}$ [®] buffer, 0.5 μl of *PrimeScript*[®] RT enzyme mix I, 0.5 μl of random 6 mers (100 μM), 0.5 μl of oligo dT primer (50 μM), and 2 μl of total RNA. The reaction programme was as follows: 37° for 15 min and 85° for 5 s. The PCR reaction mixture (25 μl) contained 12.5 μl of SYBR[®] Premix Ex TaqTM II (*TaKaRa*), 0.5 μl of PCR forward primer (10 μM), 0.5 μl of PCR reverse primer (10 μM), 9.5 μl of dH_2O , and 2 μl of cDNA. The reactions were carried out in an *iQ5* real-time PCR system (*Bio-Rad* Co., Ltd.). The reaction programme was as follows: one cycle at 95° for 30 s, followed by 40 cycles at 95° for 5 s, 60° for 30 s.

Statistical Analyses. All experiments were performed in three replications. Continuous variables, expressed as mean \pm s.d., were compared by using one-way ANOVA. Statistical analyses were conducted with SPSS 17.0.

REFERENCES

- [1] H. Li, P. Boyce, 'Pothos', in 'Flora of China', Eds. Z. Y. Wu, P. H. Raven, D. Y. Hong, Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis, 2010, Vol. 23, p. 6.
- [2] Editorial Board of Zhonghua Bencao, State Administration of Traditional Chinese Medicine of the P. R. China, 'Zhonghua Bencao', Shanghai Science and Technology Publishers, Shanghai, China, 2005, Vol. 8, p. 523.
- [3] G. Z. Yang, S. Zhao, Y. C. Li, *Yaoxue Xuebao* **2002**, 37, 437.
- [4] S. F. Hussain, B. Gözler, M. Shamma, T. Gözler, *Phytochemistry* **1980**, 21, 2979.
- [5] T. Yoshihara, S. Takamatsu, S. Sakamura, *Agric. Biol. Chem.* **1978**, 42, 623.

- [6] H. Zhang, S. Qiu, P. Tamez, G. T. Tan, Z. Aydogmus, N. Van Hung, N. M. Cuong, C. Angerhofer, D. D. Soejarto, J. M. Pezzuto, H. H. S. Fong, *Pharm Biol.* **2002**, *40*, 221.
- [7] X. Li, D. Zhang, M. Onda, Y. Konda, M. Iguchi, Y. Harigaya, *J. Nat. Prod.* **1990**, *53*, 657.
- [8] H. Chen, A. Hajia, Y. C. Li, *Nat. Prod. Res. Dev.* **2006**, *18*, 958.
- [9] F. P. Ferreira, D. C. R. de Oliveira, *Tetrahedron Lett.* **2010**, *51*, 6856.
- [10] R. E. Miller, M. J. McConville, I. E. Woodrow, *Phytochemistry* **2006**, *67*, 43.
- [11] G. M. Sheldrick, SHELXS-97, Program for Crystal Structure Solution, University of Göttingen, Göttingen, 1997.
- [12] S.-C. Ma, Z.-D. He, X.-L. Deng, P. P.-H. But, V. E.-C. Ooi, H.-X. Xu, S. H.-S. Lee, S.-F. Lee, *Chem. Pharm. Bull.* **2001**, *49*, 1471.
- [13] M. W. Pfaffl, *Nucleic Acids Res.* **2001**, *29*, 2003.
- [14] W. Lurchachaiwong, S. Payungporn, U. Srisatidnarakul, C. Mungkundar, A. Theamboonlers, Y. Poovorawan, *Lett. Appl. Microbiol.* **2008**, *46*, 55.
- [15] Y. Wang, R. Luo, L. Fang, D. Wang, J. Bi, H. Chen, S. Xiao, *Mol. Immunol.* **2011**, *48*, 586.
- [16] M. Ishiyama, M. Shiga, K. Sasamoto, M. Mizoguchi, P.-G. He, *Chem. Pharm. Bull.* **1993**, *41*, 1118.
- [17] H. Tominaga, M. Ishiyama, F. Ohseto, K. Sasamoto, T. Hamamoto, K. Suzuki, M. Watanabe, *Anal. Commun.* **1999**, *36*, 47.
- [18] Q.-Q. Huang, J.-L. Bi, Q.-Y. Sun, F.-M. Yang, Y.-H. Wang, G.-H. Tang, F.-W. Zhao, H. Wang, J.-J. Xu, E. J. Kennelly, C.-L. Long, G.-F. Yin, *Planta Med.* **2012**, *78*, 65.
- [19] B. F. Duan, Y. P. Shen, G. S. Yang, Y. F. Zhang, J. M. Wu, G. Duan, G. F. Yin, *Prog. Vet. Med.* **2010**, *31*, 11.

Received January 5, 2012